

## POLICY STATEMENT

Organizational Principles to Guide and Define the Child Health Care System and/or Improve the Health of all Children

American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

# Hospital Stay for Healthy Term Newborn Infants

William E. Benitz, MD, FAAP, COMMITTEE ON FETUS AND NEWBORN

### abstract

The hospital stay of the mother and her healthy term newborn infant should be long enough to allow identification of problems and to ensure that the mother is sufficiently recovered and prepared to care for herself and her newborn at home. The length of stay should be based on the unique characteristics of each mother-infant dyad, including the health of the mother, the health and stability of the newborn, the ability and confidence of the mother to care for herself and her newborn, the adequacy of support systems at home, and access to appropriate follow-up care in a medical home. Input from the mother and her obstetrical care provider should be considered before a decision to discharge a newborn is made, and all efforts should be made to keep a mother and her newborn together to ensure simultaneous discharge.

### PURPOSE

The purpose of this policy statement is to review issues related to length of stay and readmission of healthy term newborns and to identify specific criteria that should be met to ensure that discharge and subsequent follow-up are appropriate.

### BACKGROUND

The hospital stay of the mother and her healthy term newborn infant (mother-infant dyad) should be long enough to allow identification of problems and to ensure that the mother is sufficiently recovered and prepared to care for herself and her newborn at home. Many neonatal cardiopulmonary problems related to the transition from the intrauterine to the extrauterine environment usually become apparent during the first 12 hours after birth.<sup>1</sup> Other neonatal problems, such as jaundice,<sup>2,3</sup> ductal-dependent cardiac lesions,<sup>4,5</sup> and gastrointestinal obstruction,<sup>6</sup> may require a longer period of observation by skilled health care professionals.<sup>7</sup> Likewise, significant maternal complications, such as endometritis, may not become apparent during the first day after delivery.

*The average length of stay of the mother-infant dyad after delivery declined steadily from 1970 until the mid-1990s.<sup>8</sup> Early newborn*

FREE

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

[www.pediatrics.org/cgi/doi/10.1542/peds.2015-0699](http://www.pediatrics.org/cgi/doi/10.1542/peds.2015-0699)

DOI: 10.1542/peds.2015-0699

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics



discharge was implemented in the 1990s, but in response to the ensuing debate on the care and safety of mothers and their infants, most states and the US Congress enacted legislation that ensured hospital stay for up to 48 hours for a vaginal delivery and up to 96 hours after birth by cesarean delivery. Several subsequent studies have reported that the postpartum length-of-stay legislation has led to an increase in postpartum length of stay, but the impact of this increase in length of stay on the rate of neonatal readmissions has been inconsistent.<sup>8-11</sup>

### Risk of Readmission

Criteria for newborn discharge include physiologic stability, family preparedness and competence to provide newborn care at home, availability of social support, and access to the health care system and resources. An inadequate assessment by health care providers in any of these areas before discharge can place an infant at risk and may result in readmission. In several large epidemiologic studies, readmission rates were used to assess the adequacy of the newborn hospital length of stay. In these reports, readmissions after an early discharge varied from no increase to a significant increase.<sup>8,12-15</sup> However, the differences in the definition of early discharge, postdischarge follow-up and support, and the timing of readmissions make it difficult to compare the results. In some of these studies, the risk factors for readmission to identify infants who may benefit from either a longer hospital stay or close postdischarge follow-up also were evaluated. These studies identified jaundice, dehydration, and feeding difficulties as the most common reasons for readmission.<sup>16,17</sup> Other frequently reported risk factors for readmission were Asian race, primiparity, associated maternal morbidities, shorter gestation or lower birth weight, instrumented vaginal

delivery, and small size for gestational age.<sup>13,15-18</sup> Close follow-up and better coordination of postdischarge care were important factors in decreasing the readmission rates.<sup>13,17</sup>

### Readiness for Discharge

Readiness for discharge of a healthy term infant is traditionally determined by pediatric care providers after a review of the mother's and family members' ability to provide care to a newborn infant at home. However, perceptions about the degree of readiness at the time of discharge often differ among pediatric care providers, obstetrical care providers, and mothers.<sup>18</sup> Factors associated with perceived unreadiness for maternal or neonatal discharge, primarily as reported by mothers themselves, include first live birth, maternal history of chronic disease or illness after birth, in-hospital neonatal illness, intent to breastfeed, mothers with inadequate prenatal care and poor social support, and black non-Hispanic maternal race.<sup>13,18</sup> Although no specific clinical tool is currently available to evaluate mothers' or families' perception of readiness for discharge after delivery, the American Academy of Pediatrics Safe and Healthy Beginnings toolkit contains a discharge-readiness checklist that can aid clinicians with preparation of a newborn for discharge. This tool was tested by 22 clinical practice teams during the Safe and Healthy Beginnings improvement project and focuses on risk for severe hyperbilirubinemia, availability of breastfeeding support, and coordination of newborn care.<sup>19</sup> All efforts should be made to keep mothers and infants together to promote simultaneous discharge. To accomplish this, a pediatric care provider's decision to discharge a newborn should be made jointly with input from the mother, her obstetrical care provider, and other health care providers, such as nursing staff and social workers, who are involved in the care of the mother and her infant.

### RECOMMENDATIONS

The length of stay of a healthy term newborn should be based on the unique characteristics of each mother-infant dyad, including the health of the mother, the health and stability of the infant, the ability and confidence of the mother to care for her infant, the adequacy of support systems at home, and access to appropriate follow-up care. Input from the mother and her obstetrical care provider and nursing staff should be considered before a decision to discharge a newborn is made, and all efforts should be made to keep a mother and her newborn together to encourage on-demand breastfeeding and to ensure simultaneous discharge. It is recommended that the following minimum criteria be met before discharge of a term newborn, defined as an infant born between 37-0/7 and 41-6/7 weeks of gestation<sup>20</sup> after an uncomplicated pregnancy, labor, and delivery.

1. Clinical course and physical examination reveal no abnormalities that require continued hospitalization.
2. The infant's vital signs are documented as being within normal ranges, with appropriate variations based on physiologic state, and stable for the 12 hours preceding discharge. These ranges include an axillary temperature of 36.5°C to 37.4°C (97.7-99.3°F, measured properly in an open crib with appropriate clothing),<sup>21</sup> a respiratory rate below 60 per minute<sup>22</sup> and no other signs of respiratory distress, and an awake heart rate of 100 to 190 beats per minute.<sup>23</sup> Heart rates as low as 70 beats per minute while sleeping quietly, without signs of circulatory compromise and responding appropriately to activity, also are acceptable. Sustained heart rates near or above the upper end of this range may require further evaluation.



3. The infant has urinated regularly and passed at least 1 stool spontaneously.
4. The infant has completed at least 2 successful feedings. If the infant is breastfeeding, a caregiver knowledgeable in breastfeeding, latch, swallowing, and infant satiety should observe an actual feeding and document successful performance of these tasks in the medical record.<sup>24</sup> If the infant is bottle-feeding, it is documented that the newborn is able to coordinate sucking, swallowing, and breathing while feeding.
5. There is no evidence of excessive bleeding at the circumcision site for at least 2 hours.
6. The clinical significance of jaundice, if present before discharge, has been determined, and appropriate management and/or follow-up plans have been instituted as recommended in American Academy of Pediatrics clinical practice guidelines for management of hyperbilirubinemia.<sup>2</sup>
7. The infant has been adequately evaluated and monitored for sepsis on the basis of maternal risk factors and in accordance with current guidelines for management of neonates with suspected or proven early-onset sepsis.<sup>25</sup>
8. Maternal and infant laboratory tests are available and have been reviewed, including the following:
  - maternal syphilis, hepatitis B surface antigen, and HIV status; and
  - umbilical cord or newborn blood type and direct Coombs test result, if clinically indicated.<sup>2</sup>
9. Initial hepatitis B vaccine has been administered as indicated by the infant's risk status and according to the current immunization schedule.<sup>26</sup>
10. If the mother has not previously been vaccinated, she should receive tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis, adsorbed (Tdap) vaccine immediately after the infant is born. Other adolescents and adults who will have or anticipate having close contact with the infant should be encouraged to receive a single dose of Tdap if they have not previously received Tdap.<sup>27</sup> If a mother who delivers during the flu season has not been previously immunized, she also should receive an influenza vaccination.<sup>28</sup>
11. Newborn metabolic,<sup>29</sup> hearing,<sup>30,31</sup> and pulse oximetry<sup>32-34</sup> screenings have been completed per hospital protocol and state regulations. If screening metabolic tests were performed before 24 hours of milk feeding, a system for repeating the test during the follow-up visit must be in place in accordance with local or state policy.
12. The mother's knowledge, ability, and confidence to provide adequate care for her infant are documented by the fact that training and information has been received in the following areas:
  - the importance and benefits of breastfeeding for both mother and infant;
  - appropriate urination and stooling frequency for the infant;
  - umbilical cord, skin, and newborn genital care, as well as temperature assessment and measurement with a thermometer;
  - signs of illness and common infant problems, particularly jaundice;
  - infant safety, such as use of an appropriate car safety seat, supine positioning for sleeping, maintaining a smoke-free environment, and sleeping in proximity but not bed-sharing<sup>35,36</sup>; and
  - hand hygiene, especially as a way to reduce infection.
13. A car safety seat appropriate for the infant's maturity and medical condition that meets Federal Motor Vehicle Safety Standard 213 has been obtained and is available before hospital discharge, and the mother has demonstrated to trained hospital personnel appropriate infant positioning and use.
14. Family members or other support persons, including health care providers who are familiar with newborn care and are knowledgeable about lactation and the recognition of jaundice and dehydration, are available to the mother and infant after discharge.
15. A physician-directed source of continuing health care (medical home) for the mother and infant has been identified. Instructions to follow in the event of a complication or emergency have been provided. The mother should know how to reach the medical home and should have scheduled the infant's first visit, if possible, or know how to do so.
16. Family, environmental, and social risk factors have been assessed, and the mother and her other family members have been educated about safe home environment. When the following or other risk factors are present, discharge should be delayed until they are resolved or a plan to safeguard the newborn is in place. This plan may involve discussions with social services and/or state agencies, such as child protective services. These risk factors may include, but are not limited to the following:
  - untreated parental use of illicit substances or positive urine



toxicology results in the mother or newborn consistent with maternal abuse or misuse of drugs;

- history of child abuse or neglect by any anticipated care provider;
  - mental illness in a parent or another person in the home;
  - lack of social support, particularly for single, first-time mothers;
  - no fixed home;
  - history of domestic violence, particularly during this pregnancy;
  - adolescent mother, particularly if other previously listed conditions apply; or
  - barriers to adequate follow-up care for the newborn, such as lack of transportation to medical care services, lack of easy access to telephone communication, and non-English-speaking parents.
17. For newborns discharged before 48 hours after delivery, an appointment should be made for the infant to be examined by a health care practitioner within 48 hours of discharge.<sup>10,12,16,37,38</sup> If this cannot be ensured, discharge should be deferred until a mechanism for follow-up is identified. The follow-up visit can take place in a home, clinic, or hospital outpatient setting as long as the health care professional who examines the infant is competent in newborn assessment and the results of the follow-up visit are reported to the infant's primary care provider or his or her designee on the day of the visit. The purpose of the follow-up visit is to
- promote establishment of a relationship with the medical home by verifying the plan for health care maintenance, including a method for obtaining emergency services, preventive

care and immunizations, periodic evaluations and physical examinations, and necessary screenings;

- weigh the infant and assess the infant's general health, hydration, and degree of jaundice, and identify any new problems;
- review feeding patterns and technique, and encourage and support breastfeeding by observation of the adequacy of position, latch, and swallowing;
- obtain historical evidence of adequate stool and urine patterns;
- provide or make a referral for lactation support if the foregoing evaluations are not reassuring;
- assess quality of mother-infant attachment and details of infant behavior;
- reinforce maternal or family education in infant care, particularly regarding feeding and sleep position, avoidance of co-sleeping, and appropriate use of car safety seats, which should be used only for travel and not for positioning in the home;
- review results of outstanding laboratory tests, such as newborn metabolic screens, performed before discharge;
- perform screenings in accordance with state regulations and other tests that are clinically indicated, such as serum bilirubin; and
- assess for parental well-being with focus on screening for maternal postpartum depression.

## CONCLUSIONS

The timing of discharge from the hospital should be the decision of the health care provider caring for the mother and her newborn. This

decision should be made in consultation with the family and should not be based on arbitrary policies established by third-party payers. A shortened hospital stay (less than 48 hours after delivery) for healthy, term newborns can be accommodated but is not appropriate for every mother and newborn. If possible, institutions are encouraged to develop processes to prevent the necessity for early discharge of uninsured or underinsured newborn infants for purely financial reasons, however. Institutions should develop guidelines through their professional staff in collaboration with appropriate community agencies, including third-party payers, to establish hospital-stay programs for mothers and their healthy newborns. State and local public health agencies also should be involved in the oversight of existing hospital-stay programs for quality assurance and monitoring. Obstetrical care, newborn nursery care, and follow-up care should be considered independent services to be paid as separate packages and not as part of a global fee for maternity-newborn labor and delivery services. Adoption of standardized processes, such as predischarge checklists, may facilitate more uniform implementation of these recommendations across the full spectrum of health care settings where care for newborn infants is provided.

## LEAD AUTHOR

William E. Benitz, MD, FAAP

## COMMITTEE ON FETUS AND NEWBORN, 2014–2015

Kristi L. Watterberg, MD, FAAP, Chairperson  
Susan Aucott, MD FAAP  
William E. Benitz, MD, FAAP  
James J. Cummings, MD, FAAP  
Eric C. Eichenwald, MD, FAAP  
Jay Goldsmith, MD, FAAP  
Brenda B. Poindexter, MD, FAAP  
Karen Puopolo, MD, FAAP  
Dan L. Stewart, MD, FAAP  
Kasper S. Wang, MD, FAAP

## LIAISONS

CAPT Wanda D. Barfield, MD, MPH, FAAP – Centers for Disease Control and Prevention



James Goldberg, MD – American College of Obstetricians and Gynecologists  
 Thierry Lacaze, MD – Canadian Pediatric Society  
 Erin L. Keels, APRN, MS, NNP-BC – National Association of Neonatal Nurses  
 Tonse N. K. Raju, MD, DCH, FAAP – National Institutes of Health

## STAFF

Jim Couto, MA

## REFERENCES

- Desmond MM, Rudolph AJ, Phitaksphraiwan P. The transitional care nursery. A mechanism for preventive medicine in the newborn. *Pediatr Clin North Am.* 1966;13(3):651–668
- American Academy of Pediatrics Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics.* 2004;114(1):297–316
- Maisels MJ, Bhutani VK, Bogen D, Newman TB, Stark AR, Watchko JF. Hyperbilirubinemia in the newborn infant > or =35 weeks' gestation: an update with clarifications. *Pediatrics.* 2009;124(4):1193–1198
- Gentile R, Stevenson G, Dooley T, Franklin D, Kawabori I, Pearlman A. Pulsed Doppler echocardiographic determination of time of ductal closure in normal newborn infants. *J Pediatr.* 1981;98(3):443–448
- Lambert EC, Canent RV, Hohn AR. Congenital cardiac anomalies in the newborn. A review of conditions causing death or severe distress in the first month of life. *Pediatrics.* 1966;37(2):343–351
- Juang D, Snyder CL. Neonatal bowel obstruction. *Surg Clin North Am.* 2012;92(3):685–711, ix–x
- Jackson GL, Kennedy KA, Sendelbach DM, et al. Problem identification in apparently well neonates: implications for early discharge. *Clin Pediatr (Phila).* 2000;39(10):581–590
- Datar A, Sood N. Impact of postpartum hospital-stay legislation on newborn length of stay, readmission, and mortality in California. *Pediatrics.* 2006;118(1):63–72
- Madden JM, Soumerai SB, Lieu TA, Mandl KD, Zhang F, Ross-Degnan D; Health maintenance organization. Effects of a law against early postpartum discharge on newborn follow-up, adverse events, and HMO expenditures. *N Engl J Med.* 2002;347(25):2031–2038
- Meara E, Kotagal UR, Atherton HD, Lieu TA. Impact of early newborn discharge legislation and early follow-up visits on infant outcomes in a state Medicaid population. *Pediatrics.* 2004;113(6):1619–1627
- Madden JM, Soumerai SB, Lieu TA, Mandl KD, Zhang F, Ross-Degnan D. Length-of-stay policies and ascertainment of postdischarge problems in newborns. *Pediatrics.* 2004;113(1 pt 1):42–49
- Kotagal UR, Atherton HD, Eshett R, Schoettker PJ, Perlstein PH. Safety of early discharge for Medicaid newborns. *JAMA.* 1999;282(12):1150–1156
- Watt S, Sword W, Krueger P. Longer postpartum hospitalization options— who stays, who leaves, what changes? *BMC Pregnancy Childbirth.* 2005;5:13
- Grupp-Phelan J, Taylor JA, Liu LL, Davis RL. Early newborn hospital discharge and readmission for mild and severe jaundice. *Arch Pediatr Adolesc Med.* 1999;153(12):1283–1288
- Paul IM, Lehman EB, Hollenbeak CS, Maisels MJ. Preventable newborn readmissions since passage of the Newborns' and Mothers' Health Protection Act. *Pediatrics.* 2006;118(6):2349–2358
- Escobar GJ, Greene JD, Hulac P, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child.* 2005;90(2):125–131
- Danielsen B, Castles AG, Damberg CL, Gould JB. Newborn discharge timing and readmissions: California, 1992–1995. *Pediatrics.* 2000;106(1 pt 1):31–39
- Bernstein HH, Spino C, Finch S, et al. Decision-making for postpartum discharge of 4300 mothers and their healthy infants: the Life Around Newborn Discharge study. *Pediatrics.* 2007;120(2). Available at: [www.pediatrics.org/cgi/content/full/120/2/e391](http://www.pediatrics.org/cgi/content/full/120/2/e391)
- Safe and Healthy Beginnings. A resource toolkit for hospitals and physicians' offices. 2009. Available at: <https://www.aap.org/en-us/professional-resources/practice-support/quality-improvement/Quality-Improvement-Innovation-Networks/Pages/Safe-and-Healthy-Beginnings-A-Resource-Toolkit-for-Hospitals-and-Physicians-Offices.aspx?aid=2577>. Accessed March 3, 2014
- American College of Obstetricians and Gynecologists. ACOG Committee Opinion No 579: definition of term pregnancy. *Obstet Gynecol.* 2013;122(5):1139–1140
- Mayfield SR, Bhatia J, Nakamura KT, Rios GR, Bell EF. Temperature measurement in term and preterm neonates. *J Pediatr.* 1984;104(2):271–275
- Taylor WC, Watkins GM. Respiratory rate patterns in the newborn infant. *Can Med Assoc J.* 1960;83:1292–1295
- Semizel E, Oztürk B, Bostan OM, Cil E, Ediz B. The effect of age and gender on the electrocardiogram in children. *Cardiol Young.* 2008;18(1):26–40
- Hagan JF, Shaw JS, Duncan PM, eds. Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008
- Polin RA, Papile LA, Baley JE, et al; Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics.* 2012;129(5):1006–1015
- Centers for Disease Control and Prevention. Recommended immunization schedule for persons aged 0 through 18 years: United States—2014. Available at: [www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-schedule.pdf](http://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-schedule.pdf). Accessed March 10, 2014
- Centers for Disease Control and Prevention (CDC). Updated recommendations for use of tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccine (Tdap) in pregnant women—Advisory Committee on Immunization Practices (ACIP), 2012. *MMWR Morb Mortal Wkly Rep.* 2013;62(7):131–135
- Centers for Disease Control and Prevention. Prevention and control of seasonal influenza with vaccines. Recommendations of the Advisory Committee on Immunization Practices—United States, 2013–2014. *MMWR Recomm Rep.* 2013;62(RR-07):1–43
- American Academy of Pediatrics Newborn Screening Authoring